



INVESTOR IN PEOPLE

GB 04/2755

The Patent Office
Concept House
Cardiff Road

Newport
South Wales
NP10 8GG
WIPO

REC'D 21 JUL 2004


PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed

Dated 13 July 2004

**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

The
**Patent
Office**

28JUN03 081856-5 D00019
P01/7700 0.00/031511.5

The Patent Office

Cardiff Road
Newport
Gwent NP9 1RH

Request for grant of a patent

(See the notes on the back of this form. You can also
get an explanatory leaflet from the Patent Office to
help you fill in this form)

1. Your reference

P034697GB

27 JUN 2003

2. Patent application number

(The Patent Office will fill in this part)

0315111.5

3. Full name, address and postcode of the or of
each applicant (underline all surnames)

CANCER RESEARCH TECHNOLOGY LTD
61 Lincoln's Inn Fields
London
WC2A 3PX
(For further applicants please see continuation sheet)

8497927002

Patents ADP number (if you know it)

If the applicant is a corporate body, give the
country/state of its incorporation

UNITED KINGDOM

4. Title of the invention

SUBSTITUTED 5-MEMBERED RING COMPOUNDS AND THEIR
USE

5. Name of your agent (if you have one)

Carpmaels & Ransford

"Address for service" in the United Kingdom
to which all correspondence should be sent
(including the postcode)

43 Bloomsbury Square
London
WC1A 2RA

Patents ADP number (if you know it)

83001

6. If you are declaring priority from one or more
earlier patent applications, give the country
and the date of filing of the or of each of these
earlier applications and (if you know it) the or
each application number

Country

Priority application number
(if you know it)

Date of filing
(day / month / year)

7. If this application is divided or otherwise
derived from an earlier UK application,
give the number and the filing date of
the earlier application

Number of earlier application

Date of filing
(day / month / year)

8. Is a statement of inventorship and of right
to grant of a patent required in support of
this request? (Answer 'Yes' if:

- a) any applicant named in part 3 is not an inventor, or
- b) there is an inventor who is not named as an
applicant, or
- c) any named applicant is a corporate body

Yes

See note (d))

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form	1 ✓
Description	33 ✓
Claim(s)	6 ✓
Abstract	
Drawing(s)	

CF

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (*Patents Form 7/77*)

Request for preliminary examination and search (*Patents Form 9/77*) 1 /

Request for substantive examination (*Patents Form 10/77*)

Any other documents
(please specify)

11. I/We request the grant of a patent on the basis of this application.

Signature

Date

Carpmaels & Ransford 27th June 2003
Carpmaels & Ransford

12. Name and daytime telephone number of person to contact in the United Kingdom

P.N. HOWARD

020-7242 8692

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

Additional Applicant:-

**THE INSTITUTE OF CANCER RESEARCH
Royal Cancer Hospital
123 Old Brompton Road
London
SW7 3RP**

8173940002

State of Incorporation:

UNITED KINGDOM

Additional Applicant:-

**RIBOTARGETS LIMITED
Granta Park
Abington
Cambridge
CB1 6GB**

7964646002

State of Incorporation:

UNITED KINGDOM

Substituted 5-Membered Ring Compounds and Their Use

This invention relates to substituted 5-membered ring compounds (carbocyclic and heterocyclic compounds having 5 ring atoms) having HSP90 inhibitory activity, to the use of such compounds in medicine, in relation to diseases which are mediated by excessive or inappropriate HSP90 activity such as cancers, and to pharmaceutical compositions containing such compounds.

Background to the invention

Molecular chaperones maintain the appropriate folding and conformation of proteins and are crucial in regulating the balance between protein synthesis and degradation. They have been shown to be important in regulating many important cellular functions, such as cell proliferation and apoptosis (Jolly and Morimoto, 2000; Smith et al., 1998; Smith, 2001).

Heat Shock Proteins (HSPs)

Exposure of cells to a number of environmental stresses, including heat shock, alcohols, heavy metals and oxidative stress, results in the cellular accumulation of a number of chaperones, commonly known as heat shock proteins (HSPs). Induction of HSPs protects the cell against the initial stress insult, enhances recovery and leads to maintenance of a stress tolerant state. It has also become clear, however, that certain HSPs may also play a major molecular chaperone role under normal, stress-free conditions by regulating the correct folding, degradation, localization and function of a growing list of important cellular proteins.

A number of multigene families of HSPs exist, with individual gene products varying in cellular expression, function and localization. They are classified according to molecular weight, e.g., HSP70, HSP90, and HSP27.

Several diseases in humans can be acquired as a result of protein misfolding (reviewed in Tytell et al., 2001; Smith et al., 1998). Hence the development of therapies which disrupt the molecular chaperone machinery may prove to be beneficial. In some conditions (e.g., Alzheimer's disease, prion diseases

and Huntington's disease), misfolded proteins can cause protein aggregation resulting in neurodegenerative disorders. Also, misfolded proteins may result in loss of wild type protein function, leading to deregulated molecular and physiological functions in the cell.

5

HSPs have also been implicated in cancer. For example, there is evidence of differential expression of HSPs which may relate to the stage of tumour progression (Martin et al., 2000; Conroy et al., 1996; Kawanishi et al., 1999; Jameel et al., 1992; Hoang et al., 2000; Lebeau et al., 1991). As a result of the involvement of HSP90 in various critical oncogenic pathways and the discovery that certain natural products with anticancer activity are targeting this molecular chaperone, the fascinating new concept has been developed that inhibiting HSP function may be useful in the treatment of cancer. The first molecular chaperone inhibitor is currently undergoing clinical trials.

15

HSP90

HSP90 constitutes about 1-2% of total cellular protein, and is usually present in the cell as a dimer in association with one of a number of other proteins (see, e.g., Pratt, 1997). It is essential for cell viability and it exhibits dual chaperone functions (Young et al., 2001). It plays a key role in the cellular stress response by interacting with many proteins after their native conformation has been altered by various environmental stresses, such as heat shock, ensuring adequate protein folding and preventing non-specific aggregation (Smith et al., 1998). In addition, recent results suggest that HSP90 may also play a role in buffering against the effects of mutation, presumably by correcting the inappropriate folding of mutant proteins (Rutherford and Lindquist, 1998). However, HSP90 also has an important regulatory role. Under normal physiological conditions, together with its endoplasmic reticulum homologue GRP94, HSP90 plays a housekeeping role in the cell, maintaining the conformational stability and maturation of several key client proteins. These can be subdivided into three groups: (a) steroid hormone receptors, (b) Ser/Thr or tyrosine kinases (e.g., ERBB2, RAF-1, CDK4, and LCK), and (c) a collection of apparently unrelated proteins, e.g., mutant p53 and the catalytic subunit of telomerase hTERT. All of these

30

proteins play key regulatory roles in many physiological and biochemical processes in the cell. New HSP90 client proteins are continuously being identified.

- 5 The highly conserved HSP90 family in humans consists of four genes, namely the cytosolic HSP90 α and HSP90 β isoforms (Hickey et al., 1989), GRP94 in the endoplasmic reticulum (Argon et al., 1999) and HSP75/TRAP1 in the mitochondrial matrix (Felts et al., 2000). It is thought that all the family members have a similar mode of action, but bind to different client proteins
- 10 depending on their localization within the cell. For example, ERBB2 is known to be a specific client protein of GRP94 (Argon et al., 1999) and type 1 tumour necrosis factor receptor (TNFR1) and RB have both been shown to be clients of TRAP1 (Song et al., 1995; Chen et al., 1996).
- 15 HSP90 participates in a series of complex interactions with a range of client and regulatory proteins (Smith, 2001). Although the precise molecular details remain to be elucidated, biochemical and X-ray crystallographic studies (Prodromou et al., 1997; Stebbins et al., 1997) carried out over the last few years have provided increasingly detailed insights into the chaperone function
- 20 of HSP90.

Following earlier controversy on this issue, it is now clear that HSP90 is an ATP-dependent molecular chaperone (Prodromou et al, 1997), with dimerization of the nucleotide binding domains being essential for ATP

25 hydrolysis, which is in turn essential for chaperone function (Prodromou et al, 2000a). Binding of ATP results in the formation of a toroidal dimer structure in which the N terminal domains are brought into closer contact with each other resulting in a conformational switch known as the 'clamp mechanism' (Prodromou and Pearl, 2000b).

30

Known HSP90 Inhibitors

The first class of HSP90 inhibitors to be discovered was the benzoquinone ansamycin class, which includes the compounds herbimycin A and

geldanamycin. They were shown to reverse the malignant phenotype of fibroblasts transformed by the v-*Src* oncogene (Uehara et al., 1985), and subsequently to exhibit potent antitumour activity in both *in vitro* (Schulte et al., 1998) and *in vivo* animal models (Supko et al., 1995).

5

Immunoprecipitation and affinity matrix studies have shown that the major mechanism of action of geldanamycin involves binding to HSP90 (Whitesell et al., 1994; Schulte and Neckers, 1998). Moreover, X-ray crystallographic studies have shown that geldanamycin competes at the ATP binding site and inhibits the intrinsic ATPase activity of HSP90 (Prodromou et al., 1997; Panaretou et al., 1998). This in turn prevents the formation of mature multimeric HSP90 complexes capable of chaperoning client proteins. As a result, the client proteins are targeted for degradation via the ubiquitin proteasome pathway. 17-Allylamino, 17-demethoxygeldanamycin (17AAG) retains the property of HSP90 inhibition resulting in client protein depletion and antitumour activity in cell culture and xenograft models (Schulte et al, 1998; Kelland et al, 1999), but has significantly less hepatotoxicity than geldanamycin (Page et al, 1997). 17AAG is currently being evaluated in Phase I clinical trials.

20

Radicicol is a macrocyclic antibiotic shown to reverse the malignant phenotype of v-*Src* and v-*Ha-Ras* transformed fibroblasts (Kwon et al, 1992; Zhao et al, 1995). It was shown to degrade a number of signalling proteins as a consequence of HSP90 inhibition (Schulte et al., 1998). X-ray crystallographic data confirmed that radicicol also binds to the N terminal domain of HSP90 and inhibits the intrinsic ATPase activity (Roe et al., 1998). Radicicol lacks antitumour activity *in vivo* due to the unstable chemical nature of the compound.

25

Coumarin antibiotics are known to bind to bacterial DNA gyrase at an ATP binding site homologous to that of the HSP90. The coumarin, novobiocin, was shown to bind to the carboxy terminus of HSP90, i.e., at a different site to that occupied by the benzoquinone ansamycins and radicicol which bind at the N-terminus (Marcu et al., 2000b). However, this still resulted in inhibition

30

of HSP90 function and degradation of a number of HSP90-chaperoned signalling proteins (Marcu et al., 2000a). Geldanamycin cannot bind HSP90 subsequent to novobiocin; this suggests that some interaction between the N and C terminal domains must exist and is consistent with the view that both sites are important for HSP90 chaperone properties.

A purine-based HSP90 inhibitor, PU3, has been shown to result in the degradation of signalling molecules, including ERBB2, and to cause cell cycle arrest and differentiation in breast cancer cells (Chiosis et al., 2001).

HSP90 as a Therapeutic Target

Due to its involvement in regulating a number of signalling pathways that are crucially important in driving the phenotype of a tumour, and the discovery that certain bioactive natural products exert their effects via HSP90 activity, the molecular chaperone HSP90 is currently being assessed as a new target for anticancer drug development (Neckers et al., 1999).

The predominant mechanism of action of geldanamycin, 17AAG, and radicicol involves binding to HSP90 at the ATP binding site located in the N-terminal domain of the protein, leading to inhibition of the intrinsic ATPase activity of HSP90 (see, e.g., Prodromou et al., 1997; Stebbins et al., 1997; Panaretou et al., 1998).

Inhibition of HSP90 ATPase activity prevents recruitment of co-chaperones and encourages the formation of a type of HSP90 heterocomplex from which these client proteins are targeted for degradation via the ubiquitin proteasome pathway (see, e.g., Neckers et al., 1999; Kelland et al., 1999).

Treatment with HSP90 inhibitors leads to selective degradation of important proteins involved in cell proliferation, cell cycle regulation and apoptosis, processes which are fundamentally important in cancer.

Inhibition of HSP90 function has been shown to cause selective degradation of important signalling proteins involved in cell proliferation, cell cycle


regulation and apoptosis, processes which are fundamentally important and which are commonly deregulated in cancer (see, e.g., Hostein et al., 2001). An attractive rationale for developing drugs against this target for use in the clinic is that by simultaneously depleting proteins associated with the transformed phenotype, one may obtain a strong antitumour effect and achieve a therapeutic advantage against cancer versus normal cells. These events downstream of HSP90 inhibition are believed to be responsible for the antitumour activity of HSP90 inhibitors in cell culture and animal models (see, e.g., Schulte et al., 1998; Kelland et al., 1999).

Brief description of the invention

In copending patent applications nos. PCT/GB02/005778, GB0228417.2, GB0229618.4 and GB0309637.7, it has been shown that certain substituted pyrazoles and isoxazoles are inhibitors of HSP90 activity. Analogues of those compounds, wherein the pyrazole or isoxazole ring is replaced by other 5-membered ring systems are conformationally similar and expected to have HSP90 inhibitory activity. The present invention relates to the use of such analogous substituted 5-membered ring compounds as HSP90 inhibitors, for example for inhibition of cancer cell proliferation. A core 5-membered ring with aromatic substitution on one ring carbon atom are principle characterising features of the compounds with which the invention is concerned.

Detailed description of the invention


According to the present invention there is provided the use of a compound of formula (I) or a salt, N-oxide, hydrate or solvate thereof, in the preparation of a composition for inhibition of HSP90 activity:



wherein

ring A is an aromatic or non-aromatic carbocyclic or heterocyclic ring having 5 ring atoms;

R₁ is attached to a first ring atom of ring A and is a group of formula (IA):




wherein in any compatible combination


- 5 Ar¹ is an optionally substituted aryl or heteroaryl radical,
 Alk¹ and Alk² are optionally substituted divalent C₁-C₆ alkylene or C₂-C₆
 alkenylene radicals,
 p, r and s are independently 0 or 1,
 Z is -O-, -S-, -(C=O)-, -(C=S)-, -SO₂-, -C(=O)O-, -C(=O)NR^A-,
 10 -C(=S)NR^A-, -SO₂NR^A-, -NR^AC(=O)-, -NR^ASO₂- or -NR^A- wherein R^A
 is hydrogen or C₁-C₆ alkyl, and
 Q is hydrogen or an optionally substituted carbocyclic or heterocyclic
 radical;
- 15 R₂ is attached to a second ring atom of ring A, which is adjacent the first ring
 atom to which R₁ is attached, or is absent if that ring atom is a nitrogen atom
 which is double bonded to a neighbouring ring atom, and if not absent R₁ is
 hydrogen or
- 20 (i) a group of formula (IA) as defined in relation to R₁;
- (ii) a carboxamide radical; or
- (iii) a non aromatic carbocyclic or heterocyclic ring wherein a ring
 25 carbon is optionally substituted, and/or a ring nitrogen is optionally
 substituted by a group of formula $-(\text{Alk}^1)_p-(\text{Z})_r-(\text{Alk}^2)_s-\text{Q}$ wherein Q,
 Alk¹, Alk², Z, p, r and s are as defined above in relation to group (IA);
 and
- 30 R₃ is attached to a third ring atom of ring A, which is adjacent the second ring
 atom to which R₂ is attached, or is absent if that ring atom is a nitrogen atom
 which is double bonded to a neighbouring ring atom, and if not absent R₂ is
 hydrogen, optionally substituted cycloalkyl, cycloalkenyl, C₁-C₆ alkyl, C₁-C₆
 alkenyl, or C₁-C₆ alkynyl; or a carboxyl, carboxamide or carboxyl ester group,

PROVIDED THAT (a) at least one of R_2 and R_3 is present and is other than hydrogen and (b) the compound of formula (I) is not one of formula (IA) (IB), (IC) or (ID)


5




(IA)



(IB)



(IC)



(ID)

wherein R_1 , R_2 , and R_3 are as defined above, and R is hydrogen or optionally substituted C_1 - C_6 alkyl.

- 10 Many compounds and sub-classes of the class of compounds defined above in relation to formula (I) is believed to be novel, and the invention includes all novel members of that class and their salts, hydrates and solvates.

As used herein:

- 15 the term "carboxyl group" refers to a group of formula $-COOH$;

the term "carboxyl ester group" refers to a group of formula $-COOR$, wherein R is a radical actually or notionally derived from the hydroxyl compound ROH; and

20

the term "carboxamide group" refers to a group of formula $-CONR_aR_b$, wherein $-NR_aR_b$ is a primary or secondary (including cyclic) amino group actually or notionally derived from ammonia or the amine HNR_aR_b :

25

As used herein, the term " $(C_1$ - C_6)alkyl" refers to a straight or branched chain alkyl radical having from 1 to 6 carbon atoms, including for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, t-butyl, n-pentyl and n-hexyl.

As used herein the term "divalent (C₁-C₆)alkylene radical" means a saturated hydrocarbon chain having from 1 to 6 carbon atoms and two unsatisfied valences.

5

As used herein, the term "(C₁-C₆)alkenyl" refers to a straight or branched chain alkenyl radical having from 2 to 6 carbon atoms and containing at least one double bond of E or Z configuration, including for example, ethenyl and allyl.

10

As used herein the term "divalent (C₂-C₆)alkenylene radical" means a hydrocarbon chain having from 2 to 6 carbon atoms, at least one double bond, and two unsatisfied valences.

15

As used herein, the term "(C₁-C₆)alkynyl" refers to a straight or branched chain alkenyl radical having from 2 to 6 carbon atoms and containing at least one triple bond, including for example, ethynyl and prop-2-ynyl.

20

As used herein the term "cycloalkyl" refers to a saturated carbocyclic radical having from 3-8 carbon atoms and includes, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl.

25

As used herein the term "cycloalkenyl" refers to a carbocyclic radical having from 3-8 carbon atoms containing at least one double bond, and includes, for example, cyclopentenyl, cyclohexenyl, cycloheptenyl and cyclooctenyl.

30

As used herein the term "aryl" refers to a mono-, bi- or tri-cyclic carbocyclic aromatic radical. Illustrative of such radicals are phenyl, biphenyl and naphthyl.

As used herein the term "carbocyclic" refers to a cyclic radical whose ring atoms are all carbon, and includes monocyclic aryl, cycloalkyl and cycloalkenyl radicals.

As used herein the term "heteroaryl" refers to a mono-, bi- or tri-cyclic aromatic radical containing one or more heteroatoms selected from S, N and O. Illustrative of such radicals are thienyl, benzthienyl, furyl, benzfuryl, pyrrolyl, imidazolyl, benzimidazolyl, thiazolyl, benzthiazolyl, isothiazolyl, 5 benzisothiazolyl, pyrazolyl, oxazolyl, benzoxazolyl, isoxazolyl, benzisoxazolyl, isothiazolyl, triazolyl, benztriazolyl, thiadiazolyl, oxadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, indolyl and indazolyl.

As used herein the unqualified term "heterocyclyl" or "heterocyclic" includes 10 "heteroaryl" as defined above, and in particular means a mono-, bi- or tri-cyclic non-aromatic radical containing one or more heteroatoms selected from S, N and O, and to groups consisting of a monocyclic non-aromatic radical containing one or more such heteroatoms which is covalently linked to another such radical or to a monocyclic carbocyclic radical. Illustrative of such 15 radicals are pyrrolyl, furanyl, thienyl, piperidinyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, thiadiazolyl, pyrazolyl, pyridinyl, pyrrolidinyl, pyrimidinyl, morpholinyl, piperazinyl, indolyl, morpholinyl, benzfuranyl, pyranyl, isoxazolyl, benzimidazolyl, methylenedioxyphenyl, ethylenedioxyphenyl, maleimido and succinimido groups.

20


Unless otherwise specified in the context in which it occurs, the term "substituted" as applied to any moiety herein means substituted with at least one substituent selected from (C₁-C₆)alkyl, (C₁-C₆)alkoxy, hydroxy, hydroxy(C₁-C₆)alkyl, mercapto, mercapto(C₁-C₆)alkyl, (C₁-C₆)alkylthio, halo 25 (including fluoro and chloro), trifluoromethyl, trifluoromethoxy, nitro, nitrile (-CN), oxo, phenyl, -COOH, -COOR^A, -COR^A, -SO₂R^A, -CONH₂, -SO₂NH₂, -CONHR^A, -SO₂NHR^A, -CONR^AR^B, -SO₂NR^AR^B, -NH₂, -NHR^A, -NR^AR^B, -OCONH₂, -OCONHR^A, -OCONR^AR^B, -NHCOR^A, -NHCOOR^A, -NR^BCOOR^A, -NHCO₂OR^A, -NR^BSO₂OR^A, -NHCONH₂, -NR^ACONH₂, - 30 NHCONHR^B, -NR^ACONHR^B, -NHCONR^AR^B, or -NR^ACONR^AR^B wherein R^A and R^B are independently a (C₁-C₆)alkyl group. The term "optional substituent" means one of the foregoing substituent groups.

As used herein the term "salt" includes base addition, acid addition and quaternary salts. Compounds of the invention which are acidic can form salts, including pharmaceutically or veterinarily acceptable salts, with bases such as alkali metal hydroxides, e.g. sodium and potassium hydroxides; alkaline earth metal hydroxides e.g. calcium, barium and magnesium hydroxides; with organic bases e.g. N-ethyl piperidine, dibenzylamine and the like. Those compounds (I) which are basic can form salts, including pharmaceutically or veterinarily acceptable salts with inorganic acids, e.g. with hydrohalic acids such as hydrochloric or hydrobromic acids, sulphuric acid, nitric acid or phosphoric acid and the like, and with organic acids e.g. with acetic, tartaric, succinic, fumaric, maleic, malic, salicylic, citric, methanesulphonic and p-toluene sulphonic acids and the like.

Some compounds of the invention contain one or more actual or potential chiral centres because of the presence of asymmetric carbon atoms. The presence of several asymmetric carbon atoms gives rise to a number of diastereoisomers with R or S stereochemistry at each chiral centre. The invention includes all such diastereoisomers and mixtures thereof.

The Ring A

Compounds of formula (I) may include the following, wherein R_1 , R_2 and R_3 are as defined herein and X represents O or S:




Of the above structures, those wherein the 5-membered ring is aromatic are preferred, for example 1,2,4-tetrazolyl, or 1, 2, 3-triazole rings.

The radical R₁

In general, it is currently preferred that the radical Ar¹ present in the R₁ group is optionally substituted phenyl, preferably with one of the optional

- 5 substituents being a hydroxy group in position 2 relative to the point of attachment of the phenyl ring to the 5-membered ring. In other words, the group R₁ preferably has formula (IB)



- 10 wherein Alk¹, Alk², p, r, s, Z and Q are as defined above in relation to R₁, and R represents one or optional substituents. In the further discussion of R₁ which follows, this preference applies in addition to any other possibilities mentioned.

- 15 In the simplest structures with which the invention is concerned, each of p, r and s may be 0, and Q may be hydrogen, so that R₁ is optionally substituted aryl or heteroaryl. In such cases, R₁ may be, for example, optionally substituted phenyl, preferably 2-hydroxyphenyl which may be further substituted, for example by one or more of hydroxy, methyl, ethyl, methoxy, ethoxy, chloro, or bromo. Currently preferred are compounds wherein R₁ is
- 20 2,4-dihydroxyphenyl, substituted in the 5-position by hydroxy, methyl, ethyl, methoxy, ethoxy, chloro, or bromo.

- In other simple structures, p, r and s may again each be 0, and Q may be an optionally substituted carbocyclic or heterocyclic ring, for example phenyl,
- 25 cyclohexyl, pyridyl, morpholino, piperidiny, or piperazinyl ring. In such cases, Q is a direct substituent in the optionally substituted Ar¹ ring.

- In more complex structures with which the invention is concerned, one or more of p, r and s may be 1, and Q may be hydrogen or an optionally
- 30 substituted carbocyclic or heterocyclic ring. For example, p and/or s may be 1

- and r may be 0, so that Q is linked to Ar¹ by an alkylene or alkenylene radical, for example a C₁-C₃ alkylene radical, which is optionally substituted. In other cases each of p, r, and s may be 1, in which cases, Q is linked to Ar¹ by an alkylene or alkenylene radical which is interrupted by the hetero atom-containing Z radical. In still other cases, p and s may be 0 and r may be 1, in which case Q is linked to Ar¹ via the hetero atom-containing Z radical.

Specific examples of R₁ groups of the above types are present in the compounds of the Examples herein.

10

The radical R₂

- When R₂ is of type (i), i.e. a group of formula (IA), examples include phenyl, 2-, 3-, or 4-pyridyl, 2- or 3-furanyl, 2- or 3-thienyl, and thiazolyl wherein optional substituents include any of those listed above in the definition of "substituted", for example methoxy, ethoxy methylenedioxy, ethylenedioxy, fluoro, chloro, bromo, and trifluoromethyl. Currently preferred are compounds wherein R₂ is phenyl substituted in the 4 position by C₁-C₆ alkoxy such as methoxy or ethoxy, fluoro, chloro, bromo, morpholinomethyl, piperazino, N-methylpiperazino, or piperidino
- When R₂ is a carboxamide radical of type (ii) above, examples include those of formula -CONR^B(Alk)_nR^A wherein

25

Alk is a divalent alkylene, alkenylene or alkynylene radical, for example a -CH₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -CH₂CH=CH-, or -CH₂CCCH₂- radical, and the Alk radical may be optionally substituted,

n is 0 or 1,

30

R^B is hydrogen or a C₁-C₆ alkyl or C₂-C₆ alkenyl group, for example methyl, ethyl, n- or iso-propyl, or allyl,

R^A is hydroxy or optionally substituted carbocyclic, for example hydroxy and/or chloro-substituted phenyl and 3,4 methylenedioxyphenyl; or


heterocyclyl, for example pyridyl, furyl, thienyl, N-piperazinyl, or N-morpholinyl any of which heterocyclic rings may be substituted,

or R^A and R^B taken together with the nitrogen to which they are attached form an N-heterocyclic ring which may optionally contain one or more additional hetero atoms selected from O, S and N, and which may optionally be substituted on one or more ring C or N atoms, examples of such N-heterocyclic rings including morpholino, piperidinyl, piperazinyl and N-phenylpiperazinyl.


The radical R_3

R_3 may be, for example, hydrogen, methyl, ethyl, n- or iso-propyl, trifluoromethyl, hydroxyethyl or a carboxamide group $-\text{CONR}^B(\text{Alk})_n\text{R}^A$ as discussed above for R_2 . Hydrogen, methyl or a carboxamide group are presently preferred.

A particular sub-set of the compounds with which this invention is concerned consists of those of formula (IE) and (IF), and their salts, solvates and hydrates;




(IE)



(IF)

wherein each R independently represents an optional substituent and each of R^2 and R^3 independently represents a carboxamide group.

Another particular sub-set of the compounds with which this invention is concerned consists of those of formula (IG), and their salts, solvates and hydrates:



(IG)

wherein R₃ represents a carboxamide group (especially an ethylcarboxamide group CH₃CH₂NHC(=O)-), R₁₀ represents C₁-C₆ alkoxy such as methoxy or ethoxy, fluoro, chloro, bromo, morpholinomethyl, piperazino, N-methylpiperazino, piperidino, N-methylpiperazinomethyl, or piperidinomethyl, and R₁₁ represents bromo, chloro, phenyl, C₁-C₆ alkyl such as ethyl, isopropyl, isobutyl or tert-butyl, or phenyl(C₁-C₆ alkyl)- such as benzyl or phenylethyl.

Specific compounds with which the invention is concerned include those of the Examples.

Compounds with which the invention is concerned may be prepared by literature methods, such as those of the preparative Examples herein, and methods analogous thereto.

The compounds of the invention are inhibitors of HSP90 and are thus useful in the treatment of diseases which are mediated by excessive or inappropriate HSP90 activity such as cancers; viral diseases such as Hepatitis C (HCV) (Waxman, 2002); Immunosuppression such as in transplantation (Bijlmakers, 2000 and Yorgin, 2000); Anti-inflammatory diseases (Bucci, 2000) such as Rheumatoid arthritis, Asthma, MS, Type I Diabetes, Lupus, Psoriasis and Inflammatory Bowel Disease; Cystic fibrosis (Fuller, 2000); Angiogenesis-related diseases (Hur, 2002 and Kurebayashi, 2001): diabetic retinopathy, haemangiomas, psoriasis, endometriosis and tumour angiogenesis. Also an Hsp90 inhibitor of the invention may protect normal cells against chemotherapy-induced toxicity and be useful in diseases where failure to undergo apoptosis is an underlying factor. Such an Hsp90 inhibitor may also be useful in diseases where the induction of a cell stress or heat shock protein

response could be beneficial, for example, protection from hypoxia-ischemic injury due to elevation of Hsp70 in the heart (Hutter, 1996 and Trost, 1998) and brain (Plumier, 1997 and Rajder, 2000). An Hsp90 inhibitor could also be useful in diseases where protein misfolding or aggregation is a major causal factor, for example, scrapie/CJD, Huntingdon's and Alzheimer's (Sittler, 2001; Trazelt, 1995 and Winklhofer, 2001).

Accordingly, the invention also provides:

(i) a method of treatment of diseases or conditions mediated by excessive or inappropriate HSP90 activity in mammals, particularly humans, which method comprises administering to the mammal an amount of a compound of formula (I) as defined above, or a salt, hydrate or solvate thereof, effective to inhibit said HSP90 activity.; and

(ii) a compound of formula (I) as defined above, or a salt hydrate or solvate thereof, for use in human or veterinary medicine, particularly in the treatment of diseases or conditions mediated by excessive or inappropriate HSP90 activity;

It will be understood that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion, drug combination and the causative mechanism and severity of the particular disease undergoing therapy. In general, a suitable dose for orally administrable formulations will usually be in the range of 0.1 to 3000 mg once, twice or three times per day, or the equivalent daily amount administered by infusion or other routes. However, optimum dose levels and frequency of dosing will be determined by clinical trials as is conventional in the art.

The compounds with which the invention is concerned may be prepared for administration by any route consistent with their pharmacokinetic properties. The orally administrable compositions may be in the form of tablets, capsules,

- powders, granules, lozenges, liquid or gel preparations, such as oral, topical, or sterile parenteral solutions or suspensions. Tablets and capsules for oral administration may be in unit dose presentation form, and may contain conventional excipients such as binding agents, for example syrup, acacia, gelatin, sorbitol, tragacanth, or polyvinyl-pyrrolidone; fillers for example lactose, sugar, maize-starch, calcium phosphate, sorbitol or glycine; tableting lubricant, for example magnesium stearate, talc, polyethylene glycol or silica; disintegrants for example potato starch, or acceptable wetting agents such as sodium lauryl sulphate. The tablets may be coated according to methods well known in normal pharmaceutical practice. Oral liquid preparations may be in the form of, for example, aqueous or oily suspensions, solutions, emulsions, syrups or elixirs, or may be presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, for example sorbitol, syrup, methyl cellulose, glucose syrup, gelatin hydrogenated edible fats; emulsifying agents, for example lecithin, sorbitan monooleate, or acacia; non-aqueous vehicles (which may include edible oils), for example almond oil, fractionated coconut oil, oily esters such as glycerine, propylene glycol, or ethyl alcohol; preservatives, for example methyl or propyl p-hydroxybenzoate or sorbic acid, and if desired conventional flavouring or colouring agents.


For topical application to the skin, the drug may be made up into a cream, lotion or ointment. Cream or ointment formulations which may be used for the drug are conventional formulations well known in the art, for example as described in standard textbooks of pharmaceuticals such as the British Pharmacopoeia.

The active ingredient may also be administered parenterally in a sterile medium. Depending on the vehicle and concentration used, the drug can either be suspended or dissolved in the vehicle. Advantageously, adjuvants such as a local anaesthetic, preservative and buffering agents can be dissolved in the vehicle.

The following examples illustrate the preparation and activities of specific compounds of the invention.

Example 1


5 **1-(2,4-Dihydroxy-phenyl)-5-(4-fluoro-phenyl)-1H-[1,2,3]triazole-4-carboxylic acid**



Step 1

1-Azido-5-chloro-2,4-dimethoxy-benzene


10



To a solution of concentrated H_2SO_4 (5 mL) in water (15 mL), 5-chloro-2,4-dimethoxy-phenylamine (3g, 16mmol) was added, resulting a deep purple suspension. More water (15 mL) was then added and the mixture was cooled and stirred vigorously at 0°C in an ice-salt bath. A solution of sodium nitrite (1.2g, 17.4mmol) in water (5 mL) was added slowly and a brown solution was obtained. This solution was stirred at this temperature for an hour. After that, a solution of sodium azide (1.2g, 18.5mmol) in water (10 mL) was followed and the solution turned grey. This solution was then stirred for an hour. The grey precipitations (1.0g, 30%) formed were collected, washed with water and dried. Yellow solids formed in the filtrate were also collected, washed and dried. The second crop (1.47g, 43%) turned to light grey quickly.

^1H NMR (d_6 -acetone): 6.98 (1H, s); 6.89 (1H, s); 3.95 (3H, s) and 3.92 (3H, s)


Step 2

1-(5-Chloro-2,4-dimethoxy-phenyl)-5-(4-fluoro-phenyl)-1H-[1,2,3]triazole-4-carboxylic acid

1-Azido-5-chloro-2,4-dimethoxy-benzene (0.4g, 1.9mmol), 3-(4-Fluoro-phenyl)-3-oxo-propionic acid methyl ester (0.42g, 2.2mmol) and sodium ethoxide in EtOH (70mg of sodium in 15 mL EtOH) were refluxed for 3 hours. During the reaction, the solution turned brown from dark green. After that, EtOH was evaporated off and the resulting oil was dried in vacuum. Trituration with ether (15 mL) gave light brown solids, which were removed by filtration and washed with ether. The solids then re-dissolved in CHCl_3 and filtered. After evaporation of the solvent, light brown solids were obtained (0.35g, 50%).

^1H NMR (CDCl_3): 7.20 (1H, s, masked by solvent peak); 7.10 (2H, m); 6.70 (2H, m); 6.20 (1H, s); 3.78 (3H, s) and 3.20 (3H, s)
LC retention time 6.98 minutes $[\text{M}+\text{H}]^+$ 378.1 & 380.1 (3 : 1)

Step3

1-(2,4-Dihydroxy-phenyl)-5-(4-fluoro-phenyl)-1H-[1,2,3]triazole-4-carboxylic acid


1-(5-Chloro-2,4-dimethoxy-phenyl)-5-(4-fluoro-phenyl)-1H-[1,2,3]triazole-4-carboxylic acid (0.29g, 0.77mmol) was refluxed in a mixture of hydroiodic acid (5 mL), acetic acid (1 mL) and acetic anhydride (0.5 mL) for 16 hours. When cooled, water (20 mL) was added and the aqueous layer was extracted with EtOAc (2 x 15 mL). The combined organic layers were washed with saturated sodium thiosulphate solution (2 x 15 mL) and brine (20 mL), dried with NaSO₄ and filtered. After evaporation of the solvent, light brown solids were obtained (0.15g, 62%). The compound was used in the next synthetic step without further purification.

¹H NMR (d₆-acetone): 7.48 (2H, dd, J = 9.0 and 5.4 Hz); 7.23 (1H, dm); 7.12 (2H, dd, J = 9.0 and 8.8 Hz); 6.46 (1H, d, J = 0.6 Hz) and 6.44 (1H, dd, 8.5 and 2.5 Hz)

LC retention time 5.58 minutes [M+H]⁺ 316.1

Example 2

1-(2,4-Dihydroxy-phenyl)-5-(4-fluoro-phenyl)-1H-[1,2,3]triazole-4-carboxylic acid ethyl ester



1-(2,4-Dihydroxy-phenyl)-5-(4-fluoro-phenyl)-1H-[1,2,3]triazole-4-carboxylic acid (0.15g, 0.48mmol) was refluxed in EtOH (20 mL) in the presence of 5 drops of concentrated H₂SO₄ for 3 hours. After evaporation of the solvent, EtOAc (20 mL) was added. The organic layer was then washed with sat. NaHCO₃ (2 x 10 mL) and brine (20 mL), dried with NaSO₄ and filtered. After evaporation of the solvent, light brown semi-solids (93mg, 57%) were obtained and purified by prep. TLC (R_f = 0.73, EtOAc).


¹H NMR (d₆-acetone): 8.90 (1H, s, broad); 7.46 (2H, dd, J = 9.0 and 5.4 Hz); 7.20 (1H, dd, J = 8.6 and 0.8 Hz); 7.12 (2H, dd, J = 9.0 and 8.9 Hz); 6.46 (1H, d, J = 0.8 Hz); 6.44 (1H, dd, 8.6 and 2.5 Hz); 4.25 (2H, q, J = 7.0 Hz) and 1.20 (3H, t, J = 7.0 Hz)

5 LC retention time 6.58 minutes [M+H]⁺ 344.1

Example 3

1-(5-Chloro-2,4-dihydroxy-phenyl)-5-(4-fluoro-phenyl)-1H-[1,2,3]triazole-4-carboxylic acid ethyl ester


10



Method:

1-(5-Chloro-2-hydroxy-4-methoxy-phenyl)-5-(4-fluoro-phenyl)-1H-[1,2,3]triazole-4-carboxylic acid ethyl ester and 1-(5-Chloro-2,4-

15 dihydroxy-phenyl)-5-(4-fluoro-phenyl)-1H-[1,2,3]triazole-4-carboxylic acid ethyl ester



20 1-(5-Chloro-2,4-dimethoxy-phenyl)-5-(4-fluoro-phenyl)-1H-[1,2,3]triazole-4-carboxylic acid (38mg) was refluxed in HBr (48%, 1 mL) and acetic acid (1 mL) for 16 hours. When cooled, water (10 mL) was added and the aqueous layer was extracted with EtOAc (2 x 10 mL). The combined organic layers were then washed with brine (2 x 10 mL), dried with NaSO₄ and filtered. After
25 evaporation of the solvent, the crude acid mixture was dried in vacuum and

used in the next step. ¹Hnmr and LCMS of the crude mixture were analysed. EtOH (6 mL) and 3 drops of concentrated H₂SO₄ were added to the above acid mixture and refluxed for 2 hours. After evaporation of the solvent, EtOAc (10 mL) was added. The organic layer was then washed with sat. NaHCO₃ (2 x 10 mL) and brine (2 x 10 mL), dried with NaSO₄ and filtered. After evaporation of the solvent, the esters were purified by prep. TLC. The mono-methoxy compound has R_f = 0.52 whereas the dihydroxy compound has R_f = 0.35 (EtOAc : n-hexane / 3 : 2).

10 The mono-methoxy compound:

¹H NMR (d₆-acetone): 7.61 (1H, s); 7.45 (2H, dd, J = 9.0 and 5.4 Hz); 7.16 (2H, dd, J = 9.0 and 8.8 Hz); 6.72 (1H, s); 4.25 (2H, q, J = 7.0 Hz); 3.56 (3H, s); and 1.22 (3H, t, J = 7 Hz)

LC retention time 7.24 minutes [M-H]⁻ 390.2 & 392.1 (3 : 1)

15

The dihydroxy compound (Example 3)


¹Hnmr (d₆-acetone): 7.50 (2H, dd, J = 9.0 and 5.4 Hz); 7.48 (1H, s); 7.16 (2H, dd, J = 9.0 and 8.8 Hz); 6.66 (1H, s); 4.25 (2H, q, J = 7.0 Hz); and 1.22 (3H, t, J = 7 Hz)

20 LC retention time 7.10 minutes [M+H]⁺ 378.1 & 380.1(3 : 1)

The compounds of Examples 1-3 above were tested in the fluorescence poalarisation assay described below in the Assay section and the results are as shown in the following Table. Also tested were a number of commercially available compounds having structures with which the invention is concerned (Examples (i) – (v) in the Table). The suppliers of these commercially available compounds were Asinex Ltd (6, Schukinskaya Street, Moscow 123182, Russia) and Interbioscreen Ltd (Institutsky Prospect 7a, 142432 Chemogolovka, Russia).

30

Example	Structure	MH+	Hsp90 IC50	Source
(i)		309	B	Asinex
(ii)		287	B	Asinex
(iii)		336	A	Interbios creen
(iv)		300	A	Interbios creen
(v)		346	A	Interbios creen.
Example 1		350	B	
Example 2		344	B	


Example 3		378	A	
-----------	---	-----	---	--

Assay

- 5 A fluorescence polarization assay was employed for the evaluation of some of the compounds of the Examples:

Fluorescence Polarization Assay

- 10 Fluorescence polarization {also known as fluorescence anisotropy} measures the rotation of a fluorescing species in solution, where the larger molecule the more polarized the fluorescence emission. When the fluorophore is excited with polarized light, the emitted light is also polarized. The molecular size is proportional to the polarization of the fluorescence emission.
- 15 The fluorescein-labelled probe – RBT0045864-FAM –



- binds to HSP90 {full-length human, full-length yeast or N-terminal domain HSP90} and the anisotropy {rotation of the probe:protein complex} is measured.
- 20

Test compound is added to the assay plate, left to equilibrate and the anisotropy measured again. Any change in anisotropy is due to competitive binding of compound to HSP90, thereby releasing probe.

Materials

Chemicals are of the highest purity commercially available and all aqueous solutions are made up in AR water.

- 5 1) Costar 96-well black assay plate #3915
- 2) Assay buffer of (a) 100mM Tris pH7.4; (b) 20mM KCl; (c) 6mM MgCl₂.
Stored at room temperature.
- 3) BSA (bovine serum albumen) 10 mg/ml (New England Biolabs # B9001S)
- 4) 20 mM probe in 100 % DMSO stock concentration. Stored in the dark at
10 RT. Working concentration is 200 nM diluted in AR water and stored at 4 °C. Final concentration in assay 80 nM.
- 5) E. coli expressed human full-length HSP90 protein, purified >95% (see, e.g., Panaretou et al., 1998) and stored in 50µL aliquots at -80°C .

15 Protocol

- 1) Add 100µl 1x buffer to wells 11A and 12A (=FP BLNK)
- 2) Prepare assay mix – all reagents are kept on ice with a lid on the bucket as the probe is light-sensitive.

20

	i. Final Conc ⁿ	
• 1x Hsp90 FP Buffer	10 ml	1x
• BSA 10mg/ml (NEB)	5.0 µl	5 µg/ml
• Probe 200µM	4.0 µl	80 nM
• Human full-length Hsp90	6.25 µl	200 nM

25

- 3) Aliquot 100µl assay mix to all other wells
- 4) Seal plate and leave in dark at room temp for 20 minutes to equilibrate

Compound Dilution Plate – 1 x 3 dilution series

- 30 1) In a clear 96-well v-bottom plate – {# VWR 007/008/257} add 10 µl 100% DMSO to wells B1 to H11
- 2) To wells A1 to A11 add 17.5µl 100% DMSO

- 3) Add 2.5 μ l cpd to A1. This gives 2.5 mM {50x} stock cpd – assuming cpds 20 mM.
- 4) Repeat for wells A2 to A10. Control in columns 11 and 12.
- 5) Transfer 5 μ l from row A to row B- not column 12. Mix well.
- 5 6) Transfer 5 μ l from row B to row C. Mix well.
- 7) Repeat to row G.
- 8) Do not add any compound to row H – this is the 0 row.
- 9) This produces a 1x3 dilution series from 50 μ M to 0.07 μ M.
- 10) In well B12 prepare 20 μ l of 100 μ M standard compound.
- 10 11) After first incubation the assay plate is read on a Fusion™ a-FP plate reader (Packard BioScience, Pangbourne, Berkshire, UK).
- 12) After the first read, 2 μ l of diluted compound is added to each well for columns 1 to 10. In column 11 {provides standard curve} only add compound B11 – H11. Add 2 μ l of 100mM standard cpd to wells B12 –
- 15 H12 {is positive control }
- 13) The Z' factor is calculated from zero controls and positive wells. It typically gives a value of 0.7 – 0.9.

The compounds tested in the above assay were assigned to one of two activity ranges, namely A = <10 μ M; B = >10 μ M, and those assignments are reported above.

REFERENCES

- Argon Y and Simen BB. 1999 "Grp94, an ER chaperone with protein and peptide binding properties", Semin. Cell Dev. Biol., Vol. 10, pp. 495-505.
- Bijlmakers M-JJE, Marsh M. 2000 "Hsp90 is essential for the synthesis and subsequent membrane association, but not the maintenance, of the Src-kinase p56lck", Molecular Biology of the Cell, Vol. 11(5), pp. 1585-1595.
- Bucci M; Roviezzo F; Cicala C; Sessa WC, Cirino G. 2000 "Geldanamycin, an inhibitor of heat shock protein 90 (Hsp90) mediated signal transduction has anti-inflammatory effects and interacts with glucocorticoid receptor in vivo", Brit. J. Pharmacol., Vol 131(1), pp. 13-16.
- Chen C-F, Chen Y, Dai KD, Chen P-L, Riley DJ and Lee W-H. 1996 "A new member of the hsp90 family of molecular chaperones interacts with the retinoblastoma protein during mitosis and after heat shock", Mol. Cell. Biol., Vol. 16, pp. 4691-4699.
- Chiosis G, Timaul MN, Lucas B, Munster PN, Zheng FF, Sepp-Loenzino L and Rosen N. 2001 "A small molecule designed to bind to the adenine nucleotide pocket of HSP90 causes Her2 degradation and the growth arrest and differentiation of breast cancer cells", Chem. Biol., Vol. 8, pp. 289-299.
- Conroy SE and Latchman DS. 1996 "Do heat shock proteins have a role in breast cancer?", Brit. J. Cancer, Vol. 74, pp. 717-721.
- Felts SJ, Owen BAL, Nguyen P, Trepel J, Donner DB and Toft DO. 2000 "The HSP90-related protein TRAP1 is a mitochondrial protein with distinct functional properties", J. Biol. Chem., Vol. 5, pp. 3305-3312.
- Fuller W, Cuthbert AW. 2000 "Post-translational disruption of the delta F508 cystic fibrosis transmembrane conductance regulator (CFTR)-molecular Chaperone complex with geldanamycin stabilizes delta F508 CFTR in the rabbit reticulocyte lysate", J. Biol. Chem.; Vol 275(48), pp. 37462-37468.

- Hickey E, Brandon SE, Smale G, Lloyd D and Weber LA. 1999 "Sequence and regulation of a gene encoding a human 89-kilodalton heat shock protein", Mol. Cell. Biol., Vol. 9, pp. 2615-2626.
- 5 Hoang AT, Huang J, Rudra-Gonguly N, Zheng J, Powell WC, Rabindron SK, Wu C and Roy-Burman P. 2000 "A novel association between the human heat shock transcription factor I (HSF1) and prostate adenocarcinoma, Am. J. Pathol., Vol. 156, pp. 857-864.
- 10 Hostein I, Robertson D, Di Stefano F, Workman P and Clarke PA. 2001 "Inhibition of signal transduction by the HSP90 inhibitor 17-allylamino-17-demethoxygeldanamycin results in cytostasis and apoptosis", Cancer Res., Vol. 61, pp. 4003-4009.
- 15 Hur E, Kim H-H, Choi SM, Kim JH, Yim S, Kwon HJ, Choi Y, Kim DK, Lee M-O, Park H. 2002 "Reduction of hypoxia-induced transcription through the repression of hypoxia-inducible factor-1 α /aryl hydrocarbon receptor nuclear translocator DNA binding by the 90-kDa heat-shock protein inhibitor radicicol", Mol. Pharmacol., Vol 62(5), pp. 975-982.
- Hutter et al, 1996, Circulation, Vol.94, pp.1408.
- 20 Jameel A, Skilton RA, Campbell TA, Chander SK, Coombes RC and Luqmani YA. 1992 "Clinical and biological significance of HSP89a in human breast cancer", Int. J. Cancer, Vol. 50, pp. 409-415.
- Jolly C and Morimoto RI. 2000 "Role of the heat shock response and molecular chaperones in oncogenesis and cell death", J. Natl. Cancer Inst., Vol. 92, pp. 1564-1572.
- 25 Kawanishi K, Shiozaki H, Doki Y, Sakita I, Inoue M, Yano M, Tsujinata T, Shamma A and Monden M. 1999 "Prognostic significance of heat shock proteins 27 and 70 in patients with squamous cell carcinoma of the esophagus", Cancer, Vol. 85, pp. 1649-1657.
- 30 Kelland LR, Abel G, McKeage MJ, Jones M, Goddard PM, Valenti M, Murrer BA and Harrap KR. 1993 "Preclinical antitumour evaluation of bis-acetalo-amino-dichloro-cyclohexylamine platinum (IV): an orally active platinum drug", Cancer Research, Vol. 53, pp. 2581-2586.
- Kelland LR, Sharp SY, Rogers PM, Myers TG and Workman P. 1999 "DT-diaphorase expression and tumor cell sensitivity to 17-allylamino,

17-demethoxygeldanamycin, an inhibitor of heat shock protein 90", J. Natl. Cancer Inst., Vol. 91, pp. 1940-1949.

5 Kurebayashi J, Otsuki T, Kurosumi M, Soga S, Akinaga S, Sonoo, H. 2001 "A radicicol derivative, KF58333, inhibits expression of hypoxia-inducible factor-1 α and vascular endothelial growth factor, angiogenesis and growth of human breast cancer xenografts", Jap. J. Cancer Res., Vol 92(12), 1342-1351.

10 Kwon HJ, Yoshida M, Abe K, Horinouchi S and Bepple T. 1992 "Radicicol, an agent inducing the reversal of transformed phenotype of src-transformed fibroblasts, Biosci., Biotechnol., Biochem., Vol. 56, pp. 538-539.

15 Lebeau J, Le Cholony C, Prosperi MT and Goubin G. 1991 "Constitutive overexpression of 89 kDa heat shock protein gene in the HBL100 mammary cell line converted to a tumorigenic phenotype by the EJ/T24 Harvey-ras oncogene", Oncogene, Vol. 6, pp. 1125-1132.

Marcu MG, Chadli A, Bouhouche I, Catelli M and Neckers L. 2000a "The heat shock protein 90 antagonist novobiocin interacts with a previously unrecognized ATP-binding domain in the carboxyl terminus of the chaperone", J. Biol. Chem., Vol. 275, pp. 37181-37186.

20 Marcu MG, Schulte TW and Neckers L. 2000b "Novobiocin and related coumarins and depletion of heat shock protein 90-dependent signaling proteins", J. Natl. Cancer Inst., Vol. 92, pp. 242-248.

25 Martin KJ, Kritzman BM, Price LM, Koh B, Kwan CP, Zhang X, MacKay A, O'Hare MJ, Kaelin CM, Mutter GL, Pardee AB and Sager R. 2000 "Linking gene expression patterns to therapeutic groups in breast cancer", Cancer Res., Vol. 60, pp. 2232-2238.

Neckers L, Schulte TW and Momnaugh E. 1999 "Geldanamycin as a potential anti-cancer agent: its molecular target and biochemical activity", Invest. New Drugs, Vol. 17, pp. 361-373.

30 Page J, Heath J, Fulton R, Yalkowsky E, Tabibi E, Tomaszewski J, Smith A and Rodman L. 1997 "Comparison of geldanamycin (NSC-122750) and 17-allylaminogeldanamycin (NSC-330507D) toxicity in rats", Proc. Am. Assoc. Cancer Res., Vol. 38, pp. 308.


- Panaretou B, Prodromou C, Roe SM, O'Brien R, Ladbury JE, Piper PW and Pearl LH. 1998 "ATP binding and hydrolysis are essential to the function of the HSP90 molecular chaperone in vivo", EMBO J., Vol. 17, pp. 4829-4836.
- 5 Plumier et al, 1997, Cell. Stress Chap., Vol.2, pp.162
- Pratt WB. 1997 "The role of the HSP90-based chaperone system in signal transduction by nuclear receptors and receptors signalling via MAP kinase", Annu. Rev. Pharmacol. Toxicol., Vol. 37, pp. 297-326.
- Prodromou C and Pearl LH. 2000a "Structure and in vivo function of HSP90",
10 Curr. Opin. Struct. Biol., Vol. 10, pp. 46-51.
- Prodromou C, Roe SM, O'Brien R, Ladbury JE, Piper PW and Pearl LH. 1997 "Identification and structural characterization of the ATP/ADP-binding site in the HSP90 molecular chaperone", Cell, Vol. 90, pp. 65-75.
- Prodromou C, Panaretou B, Chohan S, Siligardi G, O'Brien R, Ladbury JE,
15 Roe SM, Piper PW and Pearl LH. 2000b "The ATPase cycle of HSP90 drives a molecular 'clamp' via transient dimerization of the N-terminal domains", EMBO J., Vol. 19, pp. 4383-4392.
- Rajder et al, 2000, Ann. Neurol., Vol.47, pp.782.
- Roe SM, Prodromou C, O'Brien R, Ladbury JE, Piper PW and Pearl LH. 1999
20 "Structural basis for inhibition of the HSP90 molecular chaperone by the antitumour antibiotics radicicol and geldanamycin", J. Med. Chem., Vol. 42, pp. 260-266.
- Rutherford SL and Lindquist S. 1998 "HSP90 as a capacitor for morphological evolution. Nature, Vol. 396, pp. 336-342.
- 25 Schulte TW, Akinaga S, Murakata T, Agatsuma T, Sugimoto S, Nakano H, Lee YS, Simen BB, Argon Y, Felts S, Toft DO, Neckers LM and Sharma SV. 1999 "Interaction of radicicol with members of the heat shock protein 90 family of molecular chaperones", Mol. Endocrinology, Vol. 13, pp. 1435-1448.
- 30 Schulte TW, Akinaga S, Soga S, Sullivan W, Sensgard B, Toft D and Neckers LM. 1998 "Antibiotic radicicol binds to the N-terminal domain of HSP90 and shares important biologic activities with geldanamycin", Cell Stress and Chaperones, Vol. 3, pp. 100-108.

- Schulte TW and Neckers LM. 1998 "The benzoquinone ansamycin 17-allylamino-17-deemthoxygeldanamycin binds to HSP90 and shares important biologic activities with geldanamycin", Cancer Chemother. Pharmacol., Vol. 42, pp. 273-279.
- 5 Sittler et al, 2001, Hum. Mol. Genet., Vol.10, pp.1307.
- Smith DF. 2001 "Chaperones in signal transduction", in: Molecular chaperones in the cell (P Lund, ed.; Oxford University Press, Oxford and NY), pp. 165-178.
- 10 Smith DF, Whitesell L and Katsanis E. 1998 "Molecular chaperones: Biology and prospects for pharmacological intervention", Pharmacological Reviews, Vol. 50, pp. 493-513.
- Song HY, Dunbar JD, Zhang YX, Guo D and Donner DB. 1995 "Identification of a protein with homology to hsp90 that binds the type 1 tumour necrosis factor receptor", J. Biol. Chem., Vol. 270, pp. 3574-3581.
- 15 Stebbins CE, Russo A, Schneider C, Rosen N, Hartl FU and Pavletich NP. 1997 "Crystal structure of an HSP90-geldanamycin complex: targeting of a protein chaperone by an antitumor agent", Cell, Vol. 89, pp. 239-250.
- 20 Supko JG, Hickman RL, Grever MR and Malspeis L. 1995 "Preclinical pharmacologic evaluation of geldanamycin as an antitumour agent", Cancer Chemother. Pharmacol., Vol. 36, pp. 305-315.
- Tratzelt et al, 1995, Proc. Nat. Acad. Sci., Vol. 92, pp. 2944.
- Trost et al, 1998, J. Clin. Invest., Vol.101, pp.855.
- 25 Tytell M and Hooper PL. 2001 "Heat shock proteins: new keys to the development of cytoprotective therapies", Emerging Therapeutic Targets, Vol. 5, pp. 267-287.
- Uehara U, Hori M, Takeuchi T and Umezawa H. 1986 "Phenotypic change from transformed to normal induced by benzoquinoid ansamycins accompanies inactivation of p60src in rat kidney cells infected with
- 30 Rous sarcoma virus", Mol. Cell. Biol., Vol. 6, pp. 2198-2206.
- Waxman, Lloyd H. Inhibiting hepatitis C virus processing and replication. (Merck & Co., Inc., USA). PCT Int. Appl. (2002), WO 0207761
- Winklhofer et al, 2001, J. Biol. Chem., Vol. 276, 45160.

- Whitesell L, Mimnaugh EG, De Costa B, Myers CE and Neckers LM. 1994
 "Inhibition of heat shock protein HSP90-pp60v-src heteroprotein
 complex formation by benzoquinone ansamycins: essential role for
 stress proteins in oncogenic transformation", Proc. Natl. Acad. Sci. U S
 5 A., Vol. 91, pp. 8324-8328.
- Yorgin et al. 2000 "Effects of geldanamycin, a heat-shock protein 90-binding
 agent, on T cell function and T cell nonreceptor protein tyrosine
 kinases", J. Immunol., Vol 164(6), pp. 2915-2923.
- 10 Young JC, Moarefi I and Hartl FU. 2001 "HSP90: a specialized but essential
 protein-folding tool", J. Cell. Biol., Vol. 154, pp. 267-273.
- Zhao JF, Nakano H and Sharma S. 1995 "Suppression of RAS and MOS
 transformation by radicicol", Oncogene, Vol. 11, pp. 161-173.

Claims:


1. The use of a compound of a compound of formula (I) or a salt, N-oxide, hydrate or solvate thereof, in the preparation of a composition for inhibition of HSP90 activity:



wherein

- 10 ring A is an aromatic or non-aromatic carbocyclic or heterocyclic ring having 5 ring atoms;

R_1 is attached to a first ring atom of ring A and is a group of formula (IA):



- 15 wherein in any compatible combination

Ar^1 is an optionally substituted aryl or heteroaryl radical,

Alk^1 and Alk^2 are optionally substituted divalent C_1 - C_6 alkylene or C_2 - C_6 alkenylene radicals,

p , r and s are independently 0 or 1,

- 20 Z is $-\text{O}-$, $-\text{S}-$, $-(\text{C}=\text{O})-$, $-(\text{C}=\text{S})-$, $-\text{SO}_2-$, $-\text{C}(=\text{O})\text{O}-$, $-\text{C}(=\text{O})\text{NR}^A-$, $-\text{C}(=\text{S})\text{NR}^A-$, $-\text{SO}_2\text{NR}^A-$, $-\text{NR}^A\text{C}(=\text{O})-$, $-\text{NR}^A\text{SO}_2-$ or $-\text{NR}^A-$ wherein R^A is hydrogen or C_1 - C_6 alkyl, and

Q is hydrogen or an optionally substituted carbocyclic or heterocyclic radical;

25

R_2 is attached to a second ring atom of ring A, which is adjacent the first ring atom to which R_1 is attached, or is absent if that ring atom is a nitrogen atom which is double bonded to a neighbouring ring atom, and if not absent R_1 is hydrogen or

30


- (i) a group of formula (IA) as defined in relation to R_1 ;

(ii) a carboxamide radical; or


(iii) a non aromatic carbocyclic or heterocyclic ring wherein a ring carbon is optionally substituted, and/or a ring nitrogen is optionally substituted by a group of formula $-(Alk^1)_p-(Z)_r-(Alk^2)_s-Q$ wherein Q, Alk^1 , Alk^2 , Z, p, r and s are as defined above in relation to group (IA); and

R_3 is attached to a third ring atom of ring A, which is adjacent the second ring atom to which R_2 is attached, or is absent if that ring atom is a nitrogen atom which is double bonded to a neighbouring ring atom, and if not absent R_2 is hydrogen, optionally substituted cycloalkyl, cycloalkenyl, C_1-C_6 alkyl, C_1-C_6 alkenyl, or C_1-C_6 alkynyl; or a carboxyl, carboxamide or carboxyl ester group,


PROVIDED THAT (a) at least one of R_2 and R_3 is present and is other than hydrogen and (b) the compound of formula (I) is not one of formula (IA) (IB), (IC) or (ID)



(IA)



(IB)



(IC)



(ID)

wherein R_1 , R_2 , and R_3 are as defined above, and R is hydrogen or optionally substituted C_1-C_6 alkyl.

2. The use as claimed in claim 1 wherein the group the ring A is aromatic.


3. The use as claimed in claim 1 or claim 2 wherein both R_1 and R_2 are attached to ring carbon atoms.

4. The use as claimed in claim 1 or claim 2 wherein one of R_1 and R_2 is attached to a ring carbon atom and the other to a ring nitrogen atom.

5. The use as claimed in claim 1 wherein the ring A is a 1,2,4-tetrazolyl

5 ring or a 1, 2, 3-triazole ring.

6. The use as claimed in any of the preceding claims wherein in the compound of formula (I) R_1 has formula (II):




10 wherein Alk^1 , Alk^2 , p , r , s , Z and Q are as defined above in relation to R_1 , and R represents one or more optional substituents.

7. The use as claimed in any of the preceding claims wherein in the group R_1 of the compound of formula (I) each of p , r and s is 0, and Q is hydrogen.

15

8. The use as claimed in claim 7 wherein R_1 is 2-hydroxyphenyl optionally further substituted by one or more of hydroxy, methyl, ethyl, methoxy, ethoxy, chloro, or bromo.

20 9. The use as claimed in any of claims 1 to 7 wherein in the compound of formula (I) R_1 has formula ((IIA):



wherein R represents bromo, chloro, phenyl, C_1 - C_6 alkyl or phenyl(C_1 - C_6 alkyl)-.

25

10. The use as claimed in any of claims 1 to 6 wherein in the group R_1 of the compound of formula (I) one or more of p , r and s is 1.

11. The use as claimed in claim 10 wherein p and/or s is/are 1 and r is 0.

12. The use as claimed in claim 10 wherein each of p, r, and s is 1.

13. The use as claimed in claim 10 wherein p and s are 0 and r is 1.

14. The use as claimed in any of the preceding claims wherein R_2 is phenyl, 2-, 3-, or 4-pyridyl, 2- or 3-furanyl, 2- or 3-thienyl, or thiazolyl, optionally substituted by one or more of methoxy, ethoxy, methylenedioxy, ethylenedioxy, fluoro, chloro, bromo, or trifluoromethyl.

15. The use as claimed in any of claims 1 to 13 wherein R_2 is optionally substituted phenyl.

16. The use as claimed in any of claims 1 to 13 wherein R_2 is a carboxamide radical of formula $-\text{CONR}^B(\text{Alk})_n\text{R}^A$ wherein

Alk is an optionally substituted divalent alkylene, alkenylene or alkynylene radical,

n is 0 or 1,

R^B is hydrogen or a $\text{C}_1\text{-C}_6$ alkyl or $\text{C}_2\text{-C}_6$ alkenyl group,

R^A is hydroxy or an optionally substituted carbocyclic or heterocyclic ring,

or R^A and R^B taken together with the nitrogen to which they are attached form an N-heterocyclic ring which may optionally contain one or more additional hetero atoms selected from O, S and N, and which may optionally be substituted on one or more ring C or N atoms.

17. The use as claimed claim 16 wherein

Alk is an optionally substituted $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}=\text{CH}-$, or $-\text{CH}_2\text{CCCH}_2-$ radical.

5 n is 0 or 1,

R^{B} is hydrogen, methyl, ethyl, n- or iso-propyl, or allyl,

10 R^{A} is hydroxy, hydroxy and/or chloro-substituted phenyl, 3,4 methylenedioxyphenyl, pyridyl, furyl, thienyl, N-piperazinyl, or N-morpholinyl,

or R^{A} and R^{B} taken together with the nitrogen to which they are attached form a morpholino, piperidiny, piperazinyl or N-phenylpiperazinyl ring.

15

18. The use as claimed in claim 16 wherein n is 0, R^{B} is hydrogen and R^{A} is hydroxy or an optionally substituted carbocyclic or heterocyclic ring.

20

19. The use as claimed in any of the preceding claims wherein R_3 is hydrogen, methyl, ethyl, n- or iso-propyl, trifluoromethyl, or hydroxyethyl.

20. The use as claimed in any of claims 1 to 18 wherein R_3 is a carboxamide group $-\text{CONR}^{\text{B}}(\text{Alk})_n\text{R}^{\text{A}}$ as defined in any of claims 16 to 18 in relation to R_2 .

25

21. A method of treatment of diseases or conditions mediated by excessive or inappropriate HSP90 activity in mammals which method comprises administering to the mammal an amount of a compound of formula (I) as defined in any of claims 1 to 20, or a salt, hydrate or solvate thereof, effective to inhibit said HSP90 activity.

30

22. The use as claimed in any of claims 1 to 20 or a method as claimed claim 21 for immunosuppression or the treatment of cancer; viral disease, inflammatory diseases such as rheumatoid arthritis, asthma, multiple

- sclerosis, Type I diabetes, lupus, psoriasis and inflammatory bowel disease; cystic fibrosis angiogenesis-related disease such as diabetic retinopathy, haemangiomas, and endometriosis; or for protection of normal cells against chemotherapy-induced toxicity; or diseases where failure to undergo
- 5 apoptosis is an underlying factor; or protection from hypoxia-ischemic injury due to elevation of Hsp70 in the heart and brain; scrapie/CJD, Huntingdon's and Alzheimer's disease.
23. A compound of formula (I) as defined in any of claims 1 to 20, or a salt
- 10 hydrate or solvate thereof, for use in human or veterinary medicine.
24. A compound of formula (I) as defined in any of claims 1 to 20, or a salt, solvate or hydrate thereof.
- 15 25. A pharmaceutical or veterinary composition comprising a compound as defined in any of claims 1 to 20, or a salt hydrate or solvate thereof, together with a pharmaceutically or veterinarily acceptable carrier.

PCT/GB2004/002755

